

Corrigendum

Molecular Biology Reports **29**: 79–82, 2002.

DNA supercoiling by gyrase is linked to nucleoid compaction

Rogier Stuger¹, Conrad L. Woldringh², Coen C. van der Weijden¹, Norbert O. E. Vischer², Barbara M. Bakker¹, Rob J.M. van Spanning¹, Jacky L. Snoep¹, and Hans V. Westerhoff^{1,2,3*}

¹*Molecular Cell Physiology, Free University, Amsterdam, The Netherlands;* ²*Swammerdam Institute for Life Sciences, University of Amsterdam, The Netherlands;* ³*Stellenbosch Institute for Advanced Study, South Africa*

*Correspondence to Hans V. Westerhoff, Molecular Cell Physiology, Free University, de Boelelaan 1085, NL-1081 HV Amsterdam, The Netherlands, EU. Email hw@bio.vu.nl, fax +31-20-444-7129, phone +31-20-444-7130.

A typographical error in Figure 2 (p. 81) escaped our attention. The relative volume of nucleoids at low DNA gyrase expression (second column, lower part of Figure 2A) is not 75, but 89. This does not affect the conclusions drawn from this experiment. The correct version of Figure 2A is printed.




Figure 2. A. DNA gyrase expression levels (Western blot with anti-gyrase A (gyrA, top), topoisomer distributions (middle), and relative average nucleoid volumes determined from fluorescence microscope images (bottom) in strain PJ4271 (control, c) and PJ4273 at increasing DNA gyrase induction levels (1, 2, 3).